

From Reactive to Proactive: Your Endometriosis Diagnostic & Management Toolkit


Reducing diagnostic delay through early recognition & coordinated care.

Identify and Intervene: Designed to address the critical diagnostic delay in endometriosis care, this toolkit is an actionable, evidence-based resource to empower Primary Care Providers (PCPs) and Emergency Medicine Practitioners (EMPs) to bridge the gap between symptom onset and specialized management through early recognition and empiric therapy.

Navigate Your Toolkit: Access foundational tools and resources to guide the care team through every stage of the endometriosis diagnostic process, including role-specific guidance where applicable. You can print this toolkit for physical reference or download the digital version to use the interactive checklists—simply click the white circles or squares to track your progress as you go.

Note: When printing, please select “Fit to Page” or “Scale to Fit” in your printer settings to preserve the formatting and readability of the document.

Support Your Practice: Explore more insights and resources through the CanSAGE Endometriosis Care Education Program (ECEP) with advanced education in imaging interpretation and surgical management pathways.



The Cost of Delay & Why Early Recognition Matters^{1,2}

A “normal” or negative imaging result is not a clean bill of health — diagnostic delay causes physical and psychological harm.


Endometriosis affects at least 1 in 10 girls, women, and unmeasured numbers of Two-Spirit and gender-diverse individuals, impacting millions of Canadians, yet patients often face significant diagnostic and treatment delays of 5+ years.

Recognize the Burden

Persistent, unmanaged symptoms contribute to physical impairment, psychological distress, and substantial socioeconomic costs, including lost productivity.

Reframe the Goal

Shift the clinical objective from “waiting for surgical confirmation” to initiating empiric therapy based on clinical suspicion.



Cycle of Fragmented Care^{1,2}

System-level gaps compound delays in diagnosis and treatment, resulting in lasting harm.

Onset & Normalization of Pain

Symptoms often begin before age 20. Adolescent pain is frequently minimized, resulting in missed opportunities for early intervention.

Fragmented Care


Patients cycle through the system without a long-term plan due to limited access to specialized diagnostics, lack of recognition and validation of symptoms, and insufficient shared decision-making in treatment options. This leads to challenges in the clinical recognition of complex symptoms, and the need for more integrated, collaborative care models.

The Emergency Pivot

Intolerable pain leads to ED visits. Without targeted follow-up, this becomes a cycle of repeat visits and perceived dismissal.

The Outcome

Undiagnosed endometriosis leads to progressive, multi-system complications that can be detrimental to productivity, social life, intimate relationships, and mental health.



Step 1 – Recognizing Endometriosis: Building Clinical Suspicion¹⁻⁵

Endometriosis is a systemic inflammatory disease often presenting with cyclic and acyclic symptoms and requiring a high index of suspicion for both pelvic and extrapelvic patterns.

The evaluation of a patient with suspected endometriosis begins with a complete history including a review of symptoms, related medical history, previous therapies, impact on quality of life, and goals of care.

Use the screening checklist on the next page to validate clinical suspicion.



History

When investigating possible endometriosis, screen for the following:

Core Symptoms (Cyclic or Acyclic):

Chronic pelvic pain (>3 months)

Common symptom cluster (4 “D”s):

Dysmenorrhea, dyspareunia,
dyschezia, dysuria

Infertility

System-Specific Symptoms (Often Cyclical):

Genital: Post-coital bleeding

Urinary: Dysuria, hematuria, flank pain

Gastrointestinal: Dyschezia, bloating,
diarrhea, constipation, nausea/emesis

Diaphragm/chest: Shoulder or subcostal
pain, pneumothorax, hemothorax

Skin, muscle, fascia: Painful mass with
cyclical exacerbation at site of previous
incisions, umbilicus

Nerve involvement: Sciatica

Additional Considerations:

Adolescent onset: Severe dysmenorrhea
starting at menarche, obstructive anomalies
(e.g., very light/absent menses with severe pain)

Impact on function: Frequent absenteeism
from school or work

Resistance to treatment: Pain unresponsive
to NSAIDs

Other risk factors: Family history (first-degree
relative), early menarche, short menstrual cycles,
heavy menstrual flow, and Mullerian anomalies

Note: Gastrointestinal symptoms and acyclic pain are common, particularly in adolescents.

Pre-exam Consultation

Before initiating physical assessment, utilize the trauma-informed protocol below to establish patient safety, consent, and control:

Validate and acknowledge patient experience

Assess clinical utility of a bimanual/internal exam: Contraindications: Adolescent patient, no history of penetrative sexual activity, or patient declines due to discomfort/trauma. A clinical diagnosis can be made, and empiric therapy initiated, based on history and/or imaging alone.

IF CONTRAINDICATED: Defer internal exam. Proceed to clinical diagnosis & empiric therapy.

IF INDICATED: Proceed with trauma-informed protocol below.

Establish patient autonomy and empower informed decision-making: Communicate to the patient that they are in control of their body and can pause/stop the exam and ask questions at any time. Ensure the results from any procedure or examination are discussed in full with the patient to empower informed decision-making and respect patient autonomy.

Obtain and maintain consent: Obtain ongoing, explicit consent for each step of the examination or any diagnostic procedure. Be mindful that some patients, particularly Indigenous women, girls, Two-Spirit, transgender, and gender diverse people, may feel pressured to comply with medical recommendations due to power imbalances and historical traumas, such as forced sterilization. Ensure patients clearly understand that no procedure will be completed without their full consent, and that they have the right to pause or stop at any moment.

Offer chaperone

Note: A bimanual examination should be performed after the myofascial assessment. The clinician should use the vaginal digit to palpate genital structures before depressing the abdominal wall with the external hand. Diagnostic caveat: Many patients will have a completely normal bimanual and speculum exam, including a normal posterior vaginal fornix. A normal physical exam does not rule out endometriosis.

Physical Examination

If a targeted physical examination is indicated, examine the following:

General well-being: Check vital signs in acute presentations (to rule out emergencies or other pathologies), mental status and BMI.

Myofascia/abdominal wall (superficial, deep, and low abdominal wall, including scars): Evaluate tone, tenderness, allodynia, or hyperalgesia for signs of central sensitization.

Neurological patterns (pain or sensory deficits).

Pelvic exam & suggestive clinical signs: A bimanual examination should be performed after the myofascial assessment. The clinician should use the vaginal digit to palpate genital structures before depressing the abdominal wall with the external hand.

Diagnostic caveat: Many patients will have a completely normal bimanual and speculum exam, including a normal posterior vaginal fornix. A normal physical exam does not rule out endometriosis.

The following clinical signs may suggest the presence of endometriosis:

Uterosacral tenderness

Fixed retroverted uterus

Ovarian masses

Cyclic scar swelling/pain


Nodularity along the vaginal fornices or cul-de-sac

Vaginal lesions of endometriosis

Hematuria/hematochezia

IF POSTERIOR COMPARTMENT YIELDS ABNORMAL FINDINGS: Consider a rectovaginal exam and kidney imaging to rule out rectal involvement and hydronephrosis.

- 1 Abdomen and Umbilicus:** Assess for bloating, myofascial tenderness/tone in the abdominal wall, and painful masses at the umbilicus or previous surgical scars.
- 2 Flank and Kidneys:** Assess for flank pain and potential hydronephrosis.
- 3 Pelvic Organs (Uterus and Ovaries):** Assess for tenderness, fixed mobility, size, and texture anomalies via bimanual exam.
- 4 Bladder:** Assess for dysuria, hematuria, and tenderness in the anterior compartment.
- 5 Bowel and Rectum:** Assess for dyschezia, constipation, diarrhea, and nodules in the posterior compartment or rectovaginal septum.
- 6 Vagina:** Assess for dyspareunia, post-coital bleeding, and visible lesions in the posterior vaginal fornix.
- 7 Lower Back and Legs:** Assess for sciatica and neurological patterns of pain or sensory deficits.



PCP Considerations:

Distinguish cyclic vs. acyclic pain patterns.

Screen for gastrointestinal or other pathology that may mimic IBS.

EMP Considerations:

Differentiate acute exacerbations of chronic pathology from acute surgical emergencies like adnexal torsion or appendicitis.

Next Step: Determine the appropriate diagnostic actions and therapy initiation.


Step 2 – Diagnostic Action Plan: Clinical Diagnosis Enables Action^{1-3,6,7}

A clinical diagnosis is sufficient to initiate triage and treatment.

Establish clinical diagnosis: A presumptive clinical diagnosis can be made based on history and physical findings to initiate empiric medical therapy.

Important to note:

- Do not use biomarkers for diagnosis as they are unreliable due to high false positives (CA-125 is often elevated in endometriosis and generally not indicated in pre-menopausal patients). Prioritize imaging characteristics instead; if scans reveal solid parts or vascularity—potential signs of malignancy—refer urgently to gynae-oncology and/or order advanced imaging.
- Do not wait for surgical confirmation before starting treatment. Clinical diagnosis is sufficient to initiate treatment for timely symptom relief.



Order Targeted Imaging

Order basic imaging to assess ovaries/uterus and rule out alternatives; order advanced imaging to map the disease and detect deep disease or multisystem involvement.^a

Note: Transvaginal scans may not be appropriate for all adolescents or non-sexually active patients. Transrectal or limited abdominal ultrasound may be used as alternatives.

Order a basic abdominal and pelvic ultrasound as first-line investigation for all patients presenting with pelvic pain to assess symptoms suggestive of endometriosis, identify ovarian endometriomas, and exclude non-endometriosis pelvic pathologies.^b

Note: A basic ultrasound assesses the uterus, ovaries, and rectouterine pouch for fluid or mass. A normal/negative basic ultrasound result does not rule out superficial endometriosis (lesions on the surface of the peritoneum) or deep endometriosis (lesions extending beyond the peritoneum).

IF NORMAL WITH NO SIGNS OF DEEP ENDOMETRIOSIS: Initiate empiric therapy.

IF DEEP ENDOMETRIOSIS IS SUSPECTED (e.g., basic ultrasound reveals an endometrioma, or the patient exhibits severe symptoms like deep dyspareunia, painful bowel/bladder habits, or a fixed/nodular uterus): Order an advanced abdominal and pelvic ultrasound for endometriosis and explicitly request 'assessment for deep endometriosis/sliding sign' (or MRI if advanced ultrasound expertise is unavailable) to diagnose ovarian endometriomas and deep endometriosis.^{b,c}

Note: An advanced ultrasound includes components of the basic ultrasound as well as assessment of the anterior compartment (bladder and ureters to assess for hydroureter), posterior compartment (bowel, uterosacral ligaments, parametria, vagina, rectovaginal septum, rectouterine pouch peritoneum), soft markers (ovarian mobility, site-specific tenderness), rectouterine pouch obliteration state (sliding sign), and kidneys (to assess for hydronephrosis).


IF OTHER PATHOLOGY: Treat accordingly.

^aCT scanning is not recommended as a primary investigative tool for pelvic endometriosis but may be indicated for acute abdominal pain or suspected thoracic endometriosis.

^b Detection of endometriomas with basic or advanced ultrasound has a sensitivity of 93% (95% CI 87%–99%) and specificity of 96% (95% CI 92%–99%).

^c Detection of deep endometriosis with advanced ultrasound has a sensitivity of 79% (95% CI 69%–89%) and specificity of 94% (95% CI 88%–100%).

Consult [How To Perform an Ultrasound to Diagnose Endometriosis](#) for a practical guide on basic versus advanced endometriosis ultrasound.



PCP: Longitudinal Management & Therapy

Goal: Treat symptoms early to prevent the development of central sensitization, chronic pelvic pain, and other long-term sequelae.

Before initiating treatment, confirm patient does not have immediate fertility goals:

If patient has immediate fertility goals: Refer for early specialist gynaecologic assessment (do not wait 1 year) and avoid hormonal suppression.

If patient does not have immediate fertility goals: Initiate therapy and use shared decision-making to personalize treatment.

Initiate Therapy: Hormonal suppression and NSAIDs for symptom relief.

Options include:

Continuous combined hormonal contraceptives at lowest dose for amenorrhea.

If abnormal uterine bleeding continues after 3 months: Increase estrogen.

If pain persists or side effects occur: Switch medication class.

Progestogens.

GnRH antagonists, GnRH agonists, and aromatase inhibitors may be considered if initial options are ineffective/contraindicated.

Consult [Diagnosis and management of endometriosis](#), published in CMAJ, for a summary of hormonal therapies for endometriosis including their dosage, adverse effects and special considerations, and relative costs.

Monitor & Adjust: Follow up at 3 months. If pain persists:

Trial a different hormonal class (e.g., switch from combined hormonal contraceptives to progestin).

Refer to Gynaecologist/Minimally Invasive Gynaecologic Surgery (MIGS) for surgical assessment.

Refer to Pelvic Health Physiotherapy for early and concurrent multidisciplinary management.

Consult [Pelvic Health Physiotherapy: A Guide for People With Endometriosis](#) for patient-focused information on pelvic health physiotherapy and how it can help manage symptoms of endometriosis and other pelvic conditions.



Proceed to the next page for EMP considerations.


EMP: The Emergency Room Pivot

Goal: Rule out the acute, validate the chronic, and bridge to care.

Differentiate and Triage: Rule out surgical emergencies (e.g., adnexal torsion, tubo-ovarian abscess, appendicitis, ruptured or hemorrhagic ovarian cyst) and non-endometrial causes of pelvic pain versus acute exacerbation of chronic endometriosis.

Check Extrapelvic Signs: If a patient presents with cyclical chest pain or shortness of breath, investigate for cyclical pneumothorax.


Provide a Clinical Bridge: Document 'suspected endometriosis' to validate the patient experience. Prescribe bridging analgesia and consider initiating hormonal suppression if no surgical emergency exists. Advise patients of available options to discuss with their primary care providers and ensure referral for outpatient follow-up.



Note: Access to advanced ultrasound expertise, pharmacotherapy costs, and insurance coverage vary significantly across provinces/territories and private plans. Please verify local formularies and eligibility criteria as coverage policies are subject to change. Next Step: Determine whether escalation or referral to specialist care is required.


Step 3 — Escalation and Referral^{1-3,8}

Timely referral to specialized care prevents disease progression, manages symptoms, and supports fertility preservation for interested patients.



Referral Checklist & Decision Algorithm

A referral is indicated if the patient meets any single criterion below. Referrals can be concurrent (e.g. to gynaecology, fertility, and/or pain specialists simultaneously) to minimize delays and address complex needs.



Surgical Indications

- Imaging confirms deep endometriosis (bowel, bladder, or ureter involvement)
- Persistent ovarian endometrioma (regardless of size) or complex ovarian cyst
- Hydroureter or hydronephrosis detected on imaging (once acute causes like kidney stones are ruled out)
- Requiring histologic confirmation (e.g., exclusion of malignancy or complex pathology)
- Symptomatic despite medical or surgical management

Refer to Gynaecologist/MIGS



Mental Health

- Patient exhibits signs of depression, anxiety, or isolation

Refer to Mental Health Specialist



Fertility Priority

- Patient desires pregnancy and has known (clinically or surgically diagnosed) or suspected endometriosis

Refer to Local Fertility Specialist or Gynaecologist
(consider concurrent referral if patient's main goal is conceiving and/or is of advanced age)



Refractory Pain

- Symptoms of endometriosis and other pelvic conditions are persistent and function-limiting
- Contraindications to hormonal therapy

Refer to Multidisciplinary Pain Support (e.g., Pelvic Physiotherapy, Chronic Pain Specialist)

Consult [Mental Health, Well-Being, and Quality of Life: A Guide for People With Endometriosis and Those Who Support Them](#) for patient-focused information on mental health strategies, therapy, and support.

When making a referral, ensure the following criteria are met:

Imaging and previous operative reports included.

Disease extent and symptom burden clearly documented.

Prior medical therapies summarized.

Fertility considerations noted.

Request specialist with advanced endometriosis surgical expertise. If a local specialist is unavailable, consider utilizing virtual consults, referring to a regional centre, and offering information on travel assistance programs.



See a [sample referral form](#) from the BC Women's Hospital + Health Centre for more details on relevant information when making a referral.


Endometriosis Diagnostic & Action Checklist¹⁻⁵

Use these clear, actionable steps to document and track your treatment progress and follow-up.

Step 1 — Recognizing Endometriosis

Step 2 — Diagnostic Action Plan

Step 3 — Escalation and Referral



Step 1 — Recognizing Endometriosis

Goal: Validate clinical suspicion using a trauma-informed approach.

Take Detailed History: Actively screen for chronic pelvic pain (>3 mos), dysmenorrhea, dyspareunia, dyschezia, dysuria, and infertility.

Review Systems: Check for cyclic symptoms in other systems: GI (bloating, diarrhea), Urinary (hematuria, flank pain), Neurological (sciatica), and Abdominal Wall (pain/mass at umbilicus or surgical scars).

Identify High-Risk Profiles: Note adolescent onset, significant absenteeism, resistance to NSAIDs, or family history.

Apply Trauma-Informed Protocol: Validate patient experience and establish patient safety and autonomy. Defer internal exams for adolescents or those without penetrative sexual history; rely on imaging instead.

Perform Physical Exam: If appropriate and consented, inspect vaginal mucosa, palpate abdominal myofascia/scars for tenderness, and perform bimanual exam to check for nodules, fixed mobility, or mass.



Step 2 — Diagnostic Action Plan

Goal: Establish a clinical diagnosis and initiate therapy without a surgically confirmed diagnosis, positive imaging, or abnormal physical exam findings.


Diagnose Clinically: Make a presumptive diagnosis based on history and physical findings to start treatment. Do not delay for surgical confirmation or use biomarkers (e.g., CA-125).

Order Imaging: Start with basic ultrasound to rule out other pelvic pathologies. Order advanced ultrasound (or MRI) if deep endometriosis is suspected or to map disease.
Tip: Explicitly request 'assessment for deep endometriosis/sliding sign' on the requisition.

PCP Considerations — Initiate Therapy: If pregnancy is not an immediate goal, prescribe NSAIDs for pain relief combined with hormonal suppression (e.g., continuous combined contraceptives or progestogens) to stop disease progression. **A confirmed diagnosis is not required to initiate therapy.**

EMP Considerations — Manage Acute Episodes: Rule out surgical emergencies (torsion, appendicitis). If stable, provide bridging analgesia and refer for outpatient management.

Monitor & Adjust: Follow up at 3 months. If pain persists, trial a different hormonal class or escalate care.



Take Home Point: Treatment can be initiated without a confirmed endometriosis diagnosis.

Step 3 — Escalation and Referral

Goal: Timely triage to prevent progression and protect fertility.


Refer to Gynaecologist/MIGS: Deep endometriosis, hydronephrosis/hydroureter, persistent/complex adnexal masses, or if medical management is ineffective.

Refer for Multidisciplinary Pain Support: Pain is function-limiting after 3 months of first-line therapy or if hormonal therapy is contraindicated or not tolerated.

Refer to Local Fertility Specialist or Gynaecologist: Known/suspected endometriosis with immediate fertility goals.


Refer for Mental Health Support: Signs of depression, anxiety, or isolation.

Prepare Referral Package: Ensure referral includes imaging reports, documentation of disease extent/symptom burden, summary of prior medical therapies, previous operative reports, and fertility goals.




The Future of Endometriosis Care¹⁻³

Proactive and continuous care leads to early action and definitive surgical management, significantly improving long-term outcomes.



Early Detection

Symptom onset, Validation by PCP, Clinical diagnosis.



Surgical Precision

Comprehensive surgery by a specialist rather than repetitive diagnostic laparoscopies.



Personalized, Patient-Centered Care

Empiric medical management, Advanced imaging if needed, Continual multidisciplinary support.



Fertility Preservation

Early counseling on fertility preservation for those at risk of ovarian damage who desire future pregnancy.

Note: Approximately 30–50% of patients with endometriosis may experience infertility.

Additional Evidence-Based Support

Support your practice with Canadian clinical standards.

- **Clinical Algorithms:** Access the [diagnosis and management of endometriosis](#) flowchart (Figure 2), published in CMAJ.
- **Imaging Guidance:** Learn [how to perform an ultrasound to diagnose endometriosis](#), published in AJUM.
- **Expert Resources:** Leverage [CanSAGE tools](#) developed by Canada's leading gynaecologic experts.
- **Center for Effective Practice (CEP) Resource:** [Endometriosis clinician tool](#) designed to support primary care clinicians in assessing, diagnosing and managing treatment for people with endometriosis.
- **Continuing Education:** Enroll in the [CanSAGE ECEP](#) to advance diagnostic and management expertise.
- **Support for People With Endometriosis:** [The Endometriosis Network Canada](#) (includes community support and resources on pelvic health physiotherapy and surgery guides).
- **Indigenous Health & Trauma-Informed Care:** Access resources such as the [Trauma-Informed Care Fact Toolkit](#) and the [Guide to Disclosure of Harm with Indigenous Patients and Families](#) to better understand the impacts of colonization, including forced sterilization and intergenerational trauma, and to build culturally safe, empowering healthcare environments for Indigenous communities.

CanSAGE acknowledges the members of the ECEP Advisory Committee for their contributions to this work. This toolkit was developed with support from Health Canada's Sexual and Reproductive Health Fund.



References

1. Singh SS, Allaire C, Al-Nourhji O, et al. Guideline No. 449: Diagnosis and Impact of Endometriosis – A Canadian Guideline. *J Obstet Gynaecol Can.* 2024;46(5):102450. doi:10.1016/j.jogc.2024.102450
2. Allaire C, Bedaiwy MA, Yong PJ. Diagnosis and management of endometriosis. *CMAJ.* 2023;195(10):E363–E371. doi:10.1503/cmaj.220637
3. ESHRE Endometriosis Guideline Group. Endometriosis – Guideline of European Society of Human Reproduction and Embryology. 2022. <https://www.eshre.eu/guideline/endometriosis>
4. Native Women’s Association of Canada. Transforming Our Approach to Promoting Sexual and Reproductive Health. 2024. https://nwac.ca/wp-content/uploads/2025/05/Trauma_Informed_Care_Fact_Toolkit.pdf
5. Alberta Health Services. Guide to Disclosure of Harm with Indigenous Patients and Families. 2025. <https://www.albertahealthservices.ca/assets/info/hp/ps/if-hp-ps-disclosure-harm-indigenous-pt-family-guide.pdf>
6. The Endometriosis Network Canada. Pelvic Health Physiotherapy: A Guide for People With Endometriosis. *Endometriosisnetwork.com.* <https://endometriosis-network.com/endo-hub/pelvic-health-physiotherapy-a-guide-for-people-with-endometriosis/>
7. Young SW, Jha P, Chamié L, et al. Society of Radiologists in Ultrasound Consensus on Routine Pelvic US for Endometriosis. *Radiology.* 2024;311(1):e232191. doi:10.1148/radiol.232191
8. BC Women’s Hospital + Health Centre. Centre for Pelvic Pain & Endometriosis Clinic Referral Form. <https://www.bcwomens.ca/Gynaecology-Site/Documents/Pelvic%20Pain-Endo/pelvic-pain-endometriosis-referral-form.pdf>

Help Us Enhance Endometriosis Care

This Endometriosis Digital Toolkit aims to empower primary care and emergency practitioners to bridge the gap between symptom onset and specialized management through early recognition and empiric therapy. To ensure this toolkit meets your clinical needs, please take our brief survey. Your insights directly improve patient care.



Take the survey.



Health
Canada

Santé
Canada

CanSAGE acknowledges the ECEP Advisory Committee for their contributions. This toolkit was supported by funding from Health Canada’s Sexual and Reproductive Health Fund. The views expressed do not necessarily reflect those of Health Canada.

